





The background of the entire page is a light gray surface with horizontal white lines, resembling a notebook. Scattered around the edges are various colorful school supplies: green rulers, orange protractors, yellow and orange paper clips, orange calculators (one showing '2368'), and pieces of paper with drawings like a cloud and a triangle. A central banner with a red-to-purple gradient contains the text.

Red/ Purple Class

Addition and Subtraction: Stage 5

Addition and Subtraction		Knowledge Organiser	
Key Vocabulary	Addition	Subtraction	
Add	Place Value Grid: $3274 + 5601 = 8875$ 	Place Value Grid: $35\ 727 - 6313 = 29\ 414$ 	
Total		2 ten thousands left 5 thousands – 6 thousands cannot be done. Exchange ten thousand for ten thousands becoming 15 thousands – 6 thousands = 9 thousands 7 hundreds – 3 hundreds = 4 hundreds 2 tens – 1 ten = 1 ten 7 ones – 3 ones = 4 ones	
Make			
Plus			
Sum			
More			
Altogether			
Difference			
Subtract			
Less			
Minus			
Take away	Column Method Starting with the ones, add each column in turn. Regroup tens, hundreds, thousands, ten thousands and/or as required. 		
Column addition			
Column subtraction			
Estimate			
Inverse operation			
Number facts			
Place value			
Complex			

Addition and Subtraction

Knowledge Organiser

Estimate and Approximate

Inverse Operations

Rounding to Estimate

41 635 + 7386 = 49 021

Round to ten:

41 630 + 7380 = 49 010

41 630 + 7390 = 49 020

41 640 + 7390 = 49 030

Rounding is not as accurate when both numbers are rounded up. A better estimate comes from "rounding" one down and one up.

Estimating on a Number Line

10 000

50 000



The arrow is about $\frac{3}{4}$ of the way across the line so it is 40 000.



Use the inverse to check:

53 476

32 732

20 744

To check $53\,476 - 32\,732 = 20\,744$
use $32\,732 + 20\,744 = 53\,476$

Start with a number, subtract 409 and double. I end with 6264. To find the starting number use the inverse: halve, then add 409. Half of 6264 = 3132. $3132 + 409 = 3541$. The starting number was 3541.

Multistep Problems

Using a Bar Model

The sum of two numbers is 25 567.

The difference is 1875.



Subtract 1875 from 25 567 = 23 692.
Halve 23 692 to find smaller number = 11 846.
Add 1875 to find larger number = 13 721.

£20

£3.75

£8.49

?

£12.24

£7.76







£20 is used to buy 2 books costing £3.75 and £8.49.




How much change is given?

£3.75 + £8.49 = £12.24

£20.00 - £12.24 = £7.76

Multiplication and Division: Stage 5

Multiplication and Division		Knowledge Organiser
Key Vocabulary	Factors	Prime Numbers
multiply	A factor is a number that divides into another number exactly, without leaving a remainder.  The factors of 20 are 1, 2, 4, 5, 10 and 20. The factor pairs are: 1 and 20 2 and 10 4 and 5 A common factor is a factor of 2 or more numbers. 	
groups of		
lots of		
times		
divide		
share		
remainder		
factor	Squared ² and Cubed ³ Numbers	Related Calculations
multiple	 $2^2 = 4$ $2 \times 2 = 4$ $2^3 = 8$ $2 \times 2 \times 2 = 8$  $5^2 = 25$ $5 \times 5 = 25$ $5^3 = 125$ $5 \times 5 \times 5 = 125$	$8 \times 9 = 72$ $80 \times 9 = 720$ $72 \div 9 = 8$ $720 \div 9 = 80$ $9 \times 8 = 72$ $90 \times 8 = 720$ $72 \div 8 = 9$ $720 \div 8 = 90$
product		
 visit twinkl.com		

Multiplication and Division		Knowledge Organiser
Short Multiplication		Long Multiplication
$2543 \times 7 = 17\ 801$  Remember to move any regrouped digits into the next column. After the next multiplication, add the regrouped number to the answer.		$2543 \times 67 = 170\ 381$  Before multiplying by the number in the tens column, remember to use zero as a placeholder because the 6 in 67 is 6 tens (60).
Division		Short Division
$136 \div 4 = 34$  30×4 4×4  visit twinkl.com		 $15 \div 4 = 3$ remainder 3 Remember to regroup any remainders and move them into the next column.  $28 \div 5 = 5$ remainder 3 If your calculation has a remainder, remember to record it in the answer using the letter r.

Four Operations: Stage 6

Four Operations

Knowledge Organiser

Key Vocabulary

Add
Total
Make
Plus
Sum
More
Altogether
Difference
Leave
Subtract
Difference between
Less
Minus
Take away
Mentally, Orally
Column Addition
Column Subtraction
Estimate
Inverse operation
Solve problems
Number facts
Place Value
Complex

Add and Subtract Whole Numbers

Column Method

	4	5	8	6	4
+	2	3	4	9	7
	6	9	3	6	1
		1	1	1	

Starting with the ones, add each column in turn. Regroup tens, hundreds, thousands, ten thousands as required.

	3	5	7	13	12
-		3	4	7	6
	3	2	2	6	6

Starting with the ones, subtract each column in turn. Exchange tens, hundreds, thousands and/or ten thousands as required.

Multiply up to 4-digit by 2-digit

1	8	2	
	1	5	4
×		2	6
	9	2	4
3	0	8	0
4	0	0	4
1	1		

Start with the ones.
 $154 \times 6 = 924$
 $154 \times 20 = 3080$
 $3080 + 924 = 4004$

Order of Operations

B	Brackets	$10 \times (4 + 2) = 10 \times 6 = 60$
O	Order	$5 + 2^2 = 5 + 4 = 9$
D	Division	$10 + 6 \div 2 = 10 + 3 = 13$
M	Multiplication	$10 - 4 \times 2 = 10 - 8 = 2$
A	Addition	$10 \times 4 + 7 = 40 + 7 = 47$
S	Subtraction	$10 \div 2 - 3 = 5 - 3 = 2$

twinkl

visit [twinkl.com](https://www.twinkl.com)

twinkl visit twinkl.com

Four Operations

Short Division

Start from the left.

		4	4	0	5
12	5	2	8	6	0

$5 \div 12 = 0 \text{ r}5$
 $52 \div 12 = 4 \text{ r}4$
 $48 \div 12 = 4$
 $6 \div 12 = 0 \text{ r}6$

Long Division

		1	2	0	r	3
14	1	6	8	3		
	1	4	0	0		
		2	8	3		
		2	8	0		
				3		

twinkl visit [twinkl.com](https://www.twinkl.com)

Knowledge Organiser

Common Factors

Factors of 48

1	2	3	4	6	8	12	16	24	48
---	---	---	---	---	---	----	----	----	----

Factors of 30

1	2	3	5	6	10	15	30
---	---	---	---	---	----	----	----

Common factors: 1, 2, 3, 6

Primes

A prime number has only 1 and itself as factors: 2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43

A composite number has factors other than 1 and itself.

Mental Calculations and Estimation

Order of calculations:

$50 \times 34 \times 2 = 50 \times 2 \times 34 = 100 \times 34 = 3400$

Money: £8.99 + £3.49 = £12.48

Use £9 + £3.50 = £12.50 and subtract 2p

Estimate on a number line

Subdivide line to estimate: 17

Common Multiples

Multiples of 3

3	...	18	21	24	...	39	42
---	-----	----	----	----	-----	----	----

Multiples of 7

7	14	21	28	35	42
---	----	----	----	----	----

Common multiples: 21, 42...

Squares and Cubes

Square numbers result from a number being multiplied by itself (e.g. $5 \times 5 = 25$):

1, 4, 9, 16, 25, 36, 49, 64, 81, 100

Cube numbers result from a number being multiplied by itself twice ($2 \times 2 \times 2 = 8$):

1, 8, 27, 64, 125

Reason from Known Facts

$90 \div 10 = 9$ so $90 \div 20 = 4.5$ and $90 \div 5 = 18$

$16 \times 9 = 144$ so $1.6 \times 9 = 14.4$

$4352 \div 17 = 256$

so $256 \times 18 = 4352 + 256 = 4608$

$3786 + 2850 = 6636$

so $4786 + 2850 = 7636$

and $2786 + 3850 = 6636$

and $8636 - 3786 = 4850$

twinkl visit twinkl.com

Fractions: Stage 5

Fractions		Knowledge Organiser
Key Vocabulary	Equivalent Fractions	Compare and Order Fractions
numerator	To find equivalent fractions, we multiply or divide the numerator and denominator by the same number.	We can compare and order fractions by using common denominators.
denominator		
unit fraction		
non-unit fraction		
whole		
equivalent	Mixed Numbers	Improper Fractions
mixed number	Mixed numbers contain a whole number and a fraction.	An improper fraction has a numerator which is greater than or equal to the denominator. $\frac{5}{3}$
improper fraction	Convert an Improper Fraction to a Mixed Number	Convert a Mixed Number to an Improper Fraction
simplest form	$\frac{9}{4}$ $9 \div 4 = 2 \text{ r } 1$ Divide the numerator by the denominator. This shows you the whole number and the fraction. $2 \frac{1}{4}$	Multiply the whole by the denominator to make an improper fraction. $2 \frac{5}{6} = \frac{12}{6} + \frac{5}{6} = \frac{17}{6}$ Add the fractions together.
multiple	Adding and Subtracting Fractions	
common denominator	To add or subtract fractions with denominators that are multiples of the same number, we must change one fraction to have the same denominator.	
common numerator		

Fractions		Knowledge Organiser						
Add Fractions Where the Total is Greater Than 1	Subtract from a Mixed Number							
$\frac{1}{2} + \frac{3}{4} + \frac{5}{8} = \frac{4}{8} + \frac{6}{8} + \frac{5}{8} = \frac{15}{8} = 1 \frac{7}{8}$	$1 \frac{2}{3} - \frac{2}{9} = 1 \frac{6}{9} - \frac{2}{9} = 1 \frac{4}{9}$ <table border="1"> <thead> <tr> <th>starting number</th><th>find the equivalent fraction</th><th>subtract</th></tr> </thead> <tbody> <tr> <td></td><td></td><td></td></tr> </tbody> </table>		starting number	find the equivalent fraction	subtract			
starting number	find the equivalent fraction	subtract						
Add Mixed Numbers	Subtract Two Mixed Numbers							
$1 \frac{1}{4} + \frac{3}{8} = 1 \frac{2}{8} + \frac{3}{8} = 1 \frac{5}{8}$ $1 \frac{1}{4} + \frac{3}{8} = \frac{5}{4} + \frac{3}{8} = \frac{10}{8} + \frac{3}{8} = \frac{13}{8} = 1 \frac{5}{8}$	$2 \frac{3}{4} - 1 \frac{5}{8} = 1 \frac{1}{8}$ $2 - 1 = 1$ $\frac{3}{4} - \frac{5}{8} = \frac{1}{8}$							
Multiply Unit Fractions by an Integer	Multiply Non-Unit Fractions by an Integer	Subtract from a Mixed Number - Breaking the Whole						
$\frac{1}{3} \times 5 = \frac{5}{3}$	$2 \times \frac{4}{9} = \frac{8}{9}$	$2 \frac{1}{4} - \frac{3}{8} = 2 \frac{2}{8} - \frac{3}{8} = 1 \frac{10}{8} - \frac{3}{8} = 1 \frac{7}{8}$						
Multiply Mixed Numbers by Integers	Subtract from a Mixed Number - Breaking the Whole							
Convert to an improper fraction and multiply the numerator by the integer. $2 \frac{1}{4} \times 2 = \frac{9}{4} \times 2 = \frac{18}{4} = 4 \frac{2}{4} = 4 \frac{1}{2}$ Use repeated addition. $2 \frac{1}{4} \times 2 = 2 \frac{1}{4} + 2 \frac{1}{4} = 4 \frac{2}{4} = 4 \frac{1}{2}$	$2 \frac{1}{4} - \frac{3}{8} = 2 \frac{2}{8} - \frac{3}{8} = 1 \frac{10}{8} - \frac{3}{8} = 1 \frac{7}{8}$							

Fractions: Stage 6

Fractions		Knowledge Organiser
Key Vocabulary	Simplify Fractions	Compare and Order Fractions
numerator denominator proper fraction improper fraction factor highest common multiple lowest common multiple equivalents common numerator common denominator decimal equivalent simplify simplest form mixed number whole number mixed number	Factors of 9: 1, 3, 9 Factors of 12: 1, 2, 3, 4, 6, 12 $\frac{9}{12} = \frac{3}{4}$	Use the Common Denominator Multiples of 5: 5, 10, 15 Multiples of 3: 3, 6, 9, 12, 15 $\frac{3}{5} = \frac{6}{10}$ $\frac{2}{3} = \frac{4}{6}$ $\frac{9}{15} < \frac{10}{15}$ $\frac{2}{3} = \frac{10}{15}$ Use the Common Numerator Multiples of 5: 5, 10, 15 Multiples of 10: 10, 20 $\frac{5}{8} = \frac{10}{16}$ $\frac{10}{16} < \frac{10}{13}$ $\frac{10}{13} = \frac{10}{13}$

Fractions		Knowledge Organiser
Adding and Subtracting Proper Fractions	Adding and Subtracting Mixed Numbers	
Same Denominators $\frac{4}{7} + \frac{2}{7} = \frac{6}{7}$ $\frac{8}{11} - \frac{3}{11} = \frac{5}{11}$ Different Denominators Multiples of 7: 7, 14, 21, 28, 35 Multiples of 5: 5, 10, 15, 20, 25, 30, 35 Multiples of 10: 10, 20 Multiples of 4: 4, 8, 12, 16, 20 $\frac{2}{7} = \frac{10}{35}$, $\frac{3}{5} = \frac{21}{35}$ $\frac{10}{35} + \frac{21}{35} = \frac{31}{35}$ $\frac{9}{10} = \frac{18}{20}$, $\frac{1}{4} = \frac{5}{20}$ $\frac{18}{20} - \frac{5}{20} = \frac{13}{20}$	Add or subtract the whole numbers and fractions separately. $2\frac{2}{5} + 1\frac{3}{10}$ $2 + 1 = 3$ $\frac{2}{5} + \frac{3}{10} = \frac{4}{10} + \frac{3}{10} = \frac{7}{10}$ $3 + \frac{7}{10} = 3\frac{7}{10}$ $2\frac{1}{2} - 1\frac{1}{4}$ $2 - 1 = 1$ $\frac{1}{2} - \frac{1}{4} = \frac{2}{4} - \frac{1}{4} = \frac{1}{4}$ $1 + \frac{1}{4} = 1\frac{1}{4}$ Convert the mixed numbers to improper fractions. $2\frac{2}{5} + 1\frac{3}{10}$ $2\frac{2}{5} = \frac{12}{5}$, $1\frac{3}{10} = \frac{13}{10}$ $\frac{12}{5} + \frac{13}{10} = \frac{24}{10} + \frac{13}{10} = \frac{37}{10}$ $\frac{37}{10} = 3\frac{7}{10}$ $2\frac{1}{2} - 1\frac{1}{4}$ $2\frac{1}{2} = \frac{5}{2}$, $1\frac{1}{4} = \frac{5}{4}$ $\frac{5}{2} - \frac{5}{4} = \frac{10}{4} - \frac{5}{4} = \frac{5}{4}$ $\frac{5}{4} = 1\frac{1}{4}$	
Multiplying Proper Fractions Multiplying Fractions by Fractions $\frac{1}{2} \times \frac{1}{3} = \frac{1 \times 1}{2 \times 3} = \frac{1}{6}$ Multiplying Fractions by Whole Numbers $\frac{2}{5} \times 3 = \frac{6}{5} = 1\frac{1}{5}$	Dividing Fractions by Whole Numbers $\frac{2}{5} \div 2 = \frac{1}{5}$ Multiplication and division are the inverse of one another so: $\div 2$ is the same as $\times \frac{1}{2}$ $\frac{2}{5} \times \frac{1}{2} = \frac{2}{10}$	

Place Value: Stage 6

Number and Place Value

Knowledge Organiser

Key Vocabulary

ten million

millions

thousands

hundreds

tens

ones

zero

place value

greater than

less than

order

round

rounded

negative number

partition

digit

interval

sequence

linear sequence

twinkl visit twinkl.com

Compare and Order

equals

$$26 + 38 = 8 \times 8$$

Both calculations have the value 64.

greater than

$$223\ 873 > 98\ 256$$

The number on the left has 2 hundred thousands and the number on the right has 0 hundred thousands.

less than

$$901\ 198 < 1\ 091\ 098$$

The number on the right has 1 million and the number on the left has 0 millions.

smallest

81 782

127 352

127 835

137 019



200 002

greatest

Negative Numbers

$$3 - 8 = -5$$

$$-6 + 11 = 5$$



Number and Place Value

Knowledge Organiser

Numbers to Ten Million

3 926 471

Millions	Hundred Thousands	Ten Thousands	Thousands	Hundreds	Tens	Ones
3	9	2	6	4	7	1


three million, nine hundred and twenty-six thousand, four hundred and seventy-one

3 926 471	
3 926 000	471



Round Any Number

Rounding to the nearest 1000



Rounding to the nearest 100 000



Rounding to the nearest 10 000



Rounding to the nearest 1 000 000



twinkl visit twinkl.com

Position and Direction: Stage 5

Position and Direction

Knowledge Organiser

Key Vocabulary

coordinate

quadrant

x-axis

y-axis

reflection

mirror line


translation

horizontal

vertical



y-axis.



Coordinates are a useful way to locate a position on a map or grid.

The numbers across the horizontal line of the grid are on the **x-axis**.

The numbers on the vertical line of the grid are on the **y-axis**.

We always read or write the number on the x-axis before the y-axis.

The x and y position are written in brackets with a comma.

The coordinate of the orange spot is **(3, 4)**.

To help you remember which point to read or write first, simply remember to move 'along the corridor and up the stairs'.

In other words, move on the **x-axis** and then move on the **y-axis**.



Position and Direction


Knowledge Organiser

Reflection

A shape is reflected when it is flipped over a mirror line.

The reflected image is congruent to the original. This means that the measurements of the sides and angles have not changed.


Each point of the reflected shape is the same distance from the mirror line as the original shape.



Translation

In maths, translation means moving an object on a grid. The object is moved without changing the size, turning or reflecting it.

When translating an object on a grid, it can move up or down, left or right.



Position and Direction: Stage 6

Position and Direction


Knowledge Organiser

Key Vocabulary

translate
translation
reflect
reflection
up
down
right
left
coordinates
quadrant
x-axis
y-axis
horizontal
vertical

Four Quadrants

Coordinates can use positive and negative numbers. Whether positive or negative, the x-axis coordinate is written first, followed by the y-axis coordinate.




Look at the circle. It is 3 units along the x-axis and 4 down the y-axis. Its coordinates are (3,-4).

Completing Shapes


Using the properties of a shape, a polygon can be completed on a grid.

To make a square, think of the square's properties.



All of a square's sides are the same length. If the completed sides are 2 units in length, the missing point must complete two more sides of 2 units.

To make a right-angled triangle, think of the triangle's properties.



A right-angled triangle should have three sides with one 90° angle.

Position and Direction


Knowledge Organiser

Translation

A shape is translated when it is moved without being rotated or resized. Every point of the shape moves the same distance and in the same direction.




Shape 1 has been translated 4 units left and 3 units down.




Reflections

A shape is reflected when it is flipped over a line which acts as a mirror. Every point on the original shape is the same distance from the mirror line as the same point on the reflected shape. The original triangle has been reflected in the x-axis and in the y-axis.



Missing Coordinates

Shapes can be shown on unmarked grids.



Point a is in the same position along the x-axis as (5,2) and in the same position on the y-axis as (7,9).

Point a (5,9)

Point b is in the same position on the y-axis as (10,4). Both triangles will have the same width. The width of the right-hand triangle is 3. This means that the width of the left-hand triangle is also 3.

Point b (2,4)

Properties of Shape:

Stage 5

Properties of Shape

Knowledge Organiser

Key Vocabulary

angle
right angle
acute
obtuse
reflex
protractor
horizontal
vertical
parallel
perpendicular
polygon
regular
irregular
two-dimensional
three-dimensional
flat face
curved surface
edge
curved edge
vertex
apex

Regular and Irregular Polygons

Regular	Irregular

A polygon is any two-dimensional shape formed with straight lines.

In a regular polygon, all the sides and angles are equal.

In an irregular polygon, the sides and angles are not equal.

Representations

Cube models can be drawn as 2D representations using different elevations.

A shape net is a 2D drawing of an unfolded 3D shape. When you are drawing or reasoning about shape nets, think carefully about where the edges of the faces meet.

Shape net of a tetrahedron.

Properties of 3D Shapes

Name	Surfaces		Edges		Vertices	Picture
	Flat	Curved	Flat	Curved		
cube	6	0	12	0	8	
cuboid	6	0	12	0	8	
square-based pyramid	5	0	8	0	5	
tetrahedron	4	0	6	0	4	
triangular prism	5	0	9	0	6	
pentagonal prism	7	0	15	0	10	
hexagonal prism	8	0	18	0	12	
octagonal prism	10	0	24	0	16	
octahedron	8	0	12	0	6	

twinkl visit twinkl.com







A cone has an apex. This is because a vertex is the point where two straight edges meet and a cone has no straight edges.



twinkl visit twinkl.com

Properties of Shape		Knowledge Organiser
Identifying Angles		
Acute Angles Any angle that measures less than 90° is called an acute angle.		Obtuse Angles Any angle that measures greater than 90° and less than 180° is called an obtuse angle.
Reflex Angles Any angle that measures greater than 180° is called a reflex angle.		
Angles on a straight line always total 180°.	Angles around a point always total 360°.	Measuring and Drawing Angles To measure angles, we use a protractor. Look carefully at how the numbers on the scale count from 0° to 180° in both directions.
Multiples of 90° can be used as descriptions of a turn.	$\frac{1}{4}$ turn - 90° $\frac{1}{2}$ turn - 180° $\frac{3}{4}$ turn - 270° 1 turn - 360°	
Using Properties of Rectangles		
	$6\text{cm} + 2\text{cm} = 8\text{cm}$ $7\text{cm} + 6\text{cm} = 13\text{cm}$	

Properties of Shape:

Stage 6

Properties of Shapes		Knowledge Organiser				
Key Vocabulary	Angle Types					
angle		Acute Angles Any angle that measures less than 90° is called an acute angle.		Obtuse Angles Any angle that measures greater than 90° and less than 180° is called an obtuse angle.		Reflex Angles Any angle that measures greater than 180° is called a reflex angle.
right angle	Calculating Angles				Angles in a Triangle	
acute						
obtuse	Angles on a straight line always total 180°.		Angles around a point always total 360°.		$a + b + c = 180^\circ$	
reflex					Angles in a Quadrilateral	
protractor	Opposite angles that share a vertex are equal.					
horizontal					$a + b + c + d = 360^\circ$	
vertical	90°		180°			
parallel						
perpendicular	270°		360°			
polygon	Multiples of 90° can be used as descriptions of a turn.					
regular						
irregular						
two-dimensional						
three-dimensional						
flat face						
curved surface						
edge						
curved edge						
vertex						
vertices						
apex						
radius						
diameter						
circumference						
 visit twinkl.com						

Properties of Shapes		Knowledge Organiser	
Using a Protractor		Angles in Regular Polygons	
Place the cross or circle at the point of the angle you are measuring. Read from the zero on the outer scale of your protractor. Count the degree lines carefully.		As the number of sides of a polygon increases by one, the total of the interior angles increases by 180°. When n = number of sides, this formula can be used to find the size of each angle in a regular polygon :	
Parts of Circles A circle is a 2D shape. The perimeter of a circle is called the circumference (c). The distance across the circle, passing through the centre, is called the diameter (d). The distance from the centre of the circle to the circumference is called the radius (r).		$\text{Sum of Interior Angles} = (n - 2) \times 180^\circ$ $\text{Each Angle} = \frac{(n - 2) \times 180^\circ}{n}$	
		 Pentagon $n = 5$ $(5 - 2) \times 180^\circ = 540^\circ$ $540^\circ \div 5 = 108^\circ$	 Hexagon $n = 6$ $(6 - 2) \times 180^\circ = 720^\circ$ $720^\circ \div 6 = 120^\circ$
Nets of 3D Shapes		Properties of 3D Shapes	
 A shape net shows which 2D shapes can be folded and joined to make a 3D shape. When you are drawing a net, or solving a problem involving a shape net, think carefully about where the edges of the faces meet.		3D shapes have three dimensions – length, width and depth . A polyhedron is a 3D shape with flat faces . Spheres, cylinders and cones are not polyhedrons as they have curved surfaces.	
 Cube 6 square faces 12 edges 8 vertices		 Tetrahedron 4 triangular faces 6 edges 4 vertices	 Sphere 1 curved surface 0 edges 0 vertices
 Cuboid 6 faces 12 edges 8 vertices		 Octahedron 8 faces 12 edges 6 vertices	 Triangular prism 5 faces 9 edges 6 vertices
 Square-based pyramid 5 faces 8 edges 5 vertices		 Cone 1 circular face 1 curved surface 1 curved edge 1 apex	 Cylinder 2 circular faces 1 curved surface 2 curved edges 0 vertices